

The Pappus Configuration, Theorem and Graph

Edward Anderson*

Abstract

We consider Projective Geometry's Pappus configuration, to which Pappus' Theorem applies. We work at the level of graphs, selecting various conceptual classes of nice presentations for this graph: Projectively-natural, Affinely-privileged, symmetric and Hamiltonian. We finally point forward to various natural successors to this configuration, fundamental Projective theorem and graph.




Figure 1:

* Dr.E.Anderson.Maths.Physics *at* protonmail.com . Institute for the Theory of STEM

This Article is (3): accessible to third-year undergraduates.

Cite as: E. Anderson, "The Pappus Configuration, Theorem and Graph", Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-pappus-graph/ (2026).

Date-stamps v1 07-02-2026, v2: 09-03-2026, v3 28-04-2026, v4 08-05-2026, v5 19-06-2026. Copyright of Dr E. Anderson.

1 Introducing Pappus' configuration and theorem

1.1 Pappus' Theorem in the Euclidean plane

Pappus' Theorem [2, 19, 18, 50, 45] Work in the usual Euclidean plane. Let A, B, C be a collinear triple of points (yellow in Fig 2). And let A', B', C' be another (red). Then

$$AB' \cap BA' := I \text{ and 3-cycles} \tag{1}$$

are themselves a collinear triple (green).




Figure 2:

Remark 1 Row 2 constructs this configuration \mathfrak{p} appus in the Affinely-special case in which the 2 input triples are parallel. While row 1 of Fig 2 depicts a generic case.

Proof 1 $\langle 0-3 \rangle^1$ One classical approach [23, 50, 45, 61] to proving this makes multiple uses of *Menelaus' Theorem*.²

Proof 2 One can proceed by multiple uses of [45] the basic *side-split lemma*. Alias *side-splitter*, *intercept* or *basic-proportionality lemma* or *theorem*, or one of the *theorems of Thales*.

Proof 3 Or by an area-sum method [45].

¹The current Encyclopedia [53] uses chevron brackets to denote "approximate year of study". This first chevron applies to the current subsection's proofs.

²See whichever of [20, 14, 13, 39] for an introduction and [23, 25, 33, 61] for discussions and applications.

Proof 4 Proofs 1 to 3 were all ‘classically available’. A further pre-Projective approach [45] is to make multiple uses of Hud–Ceva’s theorem in place of Menelaus’.

1.2 Pappus’ Theorem reappraised within Projective Geometry

Remark 1 This late Ancient Greek result harbours substantial Projective-Geometric significance. Though recognizing this had to await the 19th [7, 8, 9] and even 20th centuries [11, 28, 42]. In particular, it turns out to constitute one of the two main structural theorems of Projective Geometry. *Projective Geometry* [42, 21, 22, 12, 18, 26] is Geometry pared down to the study of *incidence*. At the level of planes, this is a binary relation on points-and-lines. Comprising whether points lie on lines and whether lines intersect each other.

Structure 1 An abstract Projective plane [55] is *Pappian* if Pappus’ theorem holds universally throughout it. Elsewise it is *non-Pappian*.

1.3 Projective-level styles of proof

Proofs 5-8 (3) It is straightforward to obtain a Projective proof intrinsically within³ $2-d$; see for instance [22, 18].

Or by using cross-ratios [47].

Or by composing Projective transformations [38].

Or by Projective transformation into a standard position followed by basic use of Coordinate Geometry [41].

Remark 1 Also in various senses Menelaus and Hud–Ceva are dual statements [61], so Proof 4 conceptually resembles proof 1.

Proof 9 (2-3) Algebraic approaches include use of homogeneous coordinates [45]. These are one of the most common types of Projectively-significant coordinates, analogous to Cartesian coordinates in Euclidean Geometry. This particular reference is further laced with Linear Algebra via its use of determinants. Though one can proceed instead using [45] (3) *Plücker relations*: another Projective notion.⁴

Proofs 10-11 (3). Once various more substantial Projective theorems are available, Pappus’ theorem drops out as a corollary.

For instance from *Pascal’s theorem*, by specializing its conic to a pair of intersecting lines [31].

Or from Desargues’ theorem, in the context of a finite plane, by a piece of Abstract Algebra.

1.4 Pedagogical aside

Pedagogical Remark 1 However that just 2 to 4 styles of proof for Pappus’ theorem suffice up to (3) . Say a classical proof in (2), though in particular the most pedestrian – side-splitter – could be attempted in some earlier year. And a Projective proof in (3), alongside a Linear Algebra proof to illustrate the ongoing usefulness of the Linear Pillar of Geometry. Those following Stillwell’s [42]

³A larger proportion of sources prove the other main theorem of Projective Geometry – Desargues’ – by exiting from $2-d$ into $3-d$.

⁴Brute-force Vector Algebra will also do [45] (5) .

Four Pillars of Geometry might pointedly want to extend your repertoire to a fourth Transformation Groups proof...

Pedagogical Remark 2 More proofs than this would be of particular interest when the Reader is doing 1) a graduate-level course in Projective Geometry. 2) A Ph.D. specializing in Projective Geometry. 3) Or a thesis or research project specifically on Pappus' theorem, or on some small set of Projective theorems that include Pappus'. Such as the Pappus–Desargues fundamental pair, or the Pascal–Brianchon dual pair [15, 23], which connects to Pappus' theorem as per Proof 10. With some tongue in cheek, passing from Stillwell's 4 Pillars of undergraduate Flat Geometry to a larger number of Pillars beyond, includes the following. Geometrical invariants themselves come to constitute a Pillar, with various further Pillars [64, 65] providing first principles either for obtaining these or for obtaining the transformation group. As such, the graduate with reasons to study Pappus' theorem further might do well to next look into a cross-ratios proof...

1.5 Optimal presentations of Pappus' configuration

Structure 1 We here aim to draw a particularly nice presentation of the Pappus configuration. For the next subsection's Graph Drawing and Visualization [48] call-sign application [36, 51]. The Pappus configuration affords the following Affine-Metric-a,d-Symmetric specialization. To parallel input lines with reflection symmetry about the perpendicular to these. Then the output line is itself parallel to these while preserving the reflection symmetry. We can then let this be a line of reflection symmetry as well. Drawing our parallel lines horizontally, this stacks the input points into 3 vertical pairs.

This leaves just 1 free parameter. Which we model as the *aspect ratio*

$$\alpha = \frac{\text{height}}{\text{width}}$$

of the bounding rectangle. Three special values for this are as follows.

$$\alpha = \frac{\sqrt{3}}{2}$$

forms equilateral triangles (Subfig a). A fortiori, it forms a patch of the regular tessellation [32] of the plane by equilateral triangles (subfig b). Rendering it a *tessellation-fitting presentation* [51]. Due to this, it is the Author's choice; Polster [36] previously made this choice as well.

It exhibits 6 little equilateral triangles (Subfig b again). Forming 2 double-length ones. With an overlap of 2 little ones (subfig c). Though this pattern extends, at the level of 6 little congruent triangles and 2 double-length triangles similar to these, throughout our 1-parameter family...

$$\alpha = 1$$

makes the bounding rectangle a square (Fig 3.d). On the one hand, its vertical sides are not realized, somewhat lessening its appeal. On the other hand, it is still isotropically proportioned, and 2 of its inclined lines are at 45° to the parallel lines. This special value was Coxeter's choice [17].

Finally

$$\alpha = \frac{1}{2}$$

places the other 4 inclined lines at 45° to the parallel lines (subfig e). This optimizes the number of (half-)right angles. Which is an Order-Theoretic Drawing and Visualization criterion [51]. Thereby, it is S. Sánchez' choice.




Figure 3:

1.6 Call-signs for pappus's objects




Figure 4:

Notational Remark 1 The Pappus objects highlighted in fireopal are now simplified, while all the unselected objects previously cast in jet are now jettisoned. The resulting ‘call-signs’ are presented inside circles, since our main use of them shall be to label the nodes of the Pappus incidence graph in the sequel Article [57]. This notation follows on from S. Sánchez’ devising of call-signs for the Fano objects [51, 55]; a different species of call-sign appear in Polster’s book [36]. We shall furthermore on occasion pick out the input and output objects by colouring the corresponding call-sign shapes to match Fig 2. Fano had no need of such colour-coding, but it turns out that both Pappus and Desargues do.

1.7 pappus’s symmetries

Exercise 1 These form a group of order

$$108 = 2^2 \times 3^3 .$$

Or, including the duality generator, of order

$$216 = 6^3 = 2^3 \times 3^3 .$$

Justify this count.

Pointer 1 Working out which group this is – i.e. relators as well as generators – lies beyond the scope of the current Encyclopedia. See Coxeter’s articles [29, 30] for details (6-8) .

2 Introducing the Pappus graph

2.1 The Projectively-natural presentation

Remark 1 Graphs associated with Geometrical figures are Topological-level constructs. So we can recycle the above nice equilateral-triangles subcase of $\mathfrak{p}_{\text{appus}}$ to form a nice presentation for the corresponding Pappus configuration graph, $\mathfrak{P}_{\text{appus}}$. See Fig 1.a). This is not to be confused with the aforementioned Pappus incidence graph...

2.2 Some basic counts and properties

Remark 2 So the Pappus graph has

$$V(\mathfrak{p}_{\text{appus}}) = N = 9, \quad (2)$$

$$E(\mathfrak{p}_{\text{appus}}) = 18. \quad (3)$$

Remark 3 The Pappus graph's degree-vector sequence is

$$\text{dv}(\mathfrak{p}_{\text{appus}}) = 3^4 4^2 5^2 6. \quad (4)$$

The Pappus graph is consequently neither *regular* [52]: of a single degree. Nor a *cone* [52]: with

$$\geq 1 \text{ degree-}(V - 1) = 8 \text{ vertices}.$$

It does however have a sole vertex with greater degree strength than the others: 6. Which we subsequently prefer to place centrally in forming nice presentation. And on average it has exactly 4 edges per vertex.

Remark 4 It is straightforward to show that the Pappus graph is *planar* [27, 40]. Given Fig 1.a)'s presentation 'in the wild', 4 identical tucks of the degree-3 vertices will do. The square-perimeter 30° angle presentation of this is exhibited in Subfig b).

Remark 5 The Pappus graph then clearly has

$$F(\mathfrak{p}_{\text{appus}}) = 10$$

internal faces Or a total of

$$F_{\text{T}}(\mathfrak{p}_{\text{appus}}) = 11$$

faces including the external face.

In the first interpretation,

$$V - E + F = 9 - 18 + 10 = 1 = \chi(\mathbb{D}^2) :$$

the Euler characteristic of the disc. In the second interpretation,

$$V - E + F_{\text{T}} = 9 - 18 + 11 = 2 = \chi(\mathbb{S}^2) .$$

Where \mathbb{D}^2 is the disc and \mathbb{S}^2 is the sphere.

2.3 Grid and tessellation presentations




Figure 5:

Remark 6 We provide four of these in Fig 5.

2.4 Ramsey presentations






	Equilateral-triangle subcase of Projectively-natural presentation	Square-perimeter 30° planar presentation
Graph	a) 	b) 
Ramsey presentation	c) 	d) 
Graph complement	e) 	<div style="border: 1px solid black; padding: 10px; text-align: center;"> <p>Pappus graph: with Ramsay presentation and complement.</p> <p><small>© 2026 Dr E. Anderson</small></p> </div>

Figure 6:

Structure 1 See Subfigs 6.c-d) for the corresponding *Ramsey presentations* on each of the already-encountered presentations in Subfigs a-b)'s blue chassis. This builds in a simple graph's non-edges in red: on an equal footing with the edges in blue.

Remark 7 One can then peel off the blue to leave the red to subsequently unfold, revealing the structure of one's incipient graph's complement. See row 3 for the peeled version, and Fig 7 for unfolded versions.

2.5 The Pappus graph is edge-critical and heterocomplementary

Structure 2 A simple graph G is *edge-critical* if

$$E(G; N) = E_{\text{crit}}(N) = \frac{V(V-1)}{4}. \quad (5)$$

This corresponds to a graph G and its complement \overline{G} having the same number of edges. This can only happen if

$$V = 0, 1 \pmod{4}. \quad (6)$$

Structure 3 A graph is *homocomplementary* if it is isomorphic to its own complement. Otherwise it is *heterocomplementary*.⁵

Diagnostic 1 The above simple graph edge-criticality conditions (5, 6) are necessary but not sufficient for a graph to be homocomplementary.

Remark 8 In fact, generically, edge-critical simple graphs are not self-complementary.

Diagnostic 2 A more detailed necessary but not sufficient condition for homocomplementarity is that the graph and its complement share degree sequence.

Remark 9 From (2, 3), the Pappus graph turns out to be edge-critical:

$$E_{\text{crit}}(9) = \frac{V(V-1)}{4} = \frac{9(9-1)}{4} = \frac{9 \times 8}{4} = 18 = E(\mathfrak{p}_{\text{appus}}; 9).$$

And yet to be generic in the heterocomplementary sense rather than homocomplementary. For

$$\deg(\overline{\mathfrak{p}_{\text{appus}}}) = 2 \cdot 3^2 \cdot 4^2 \cdot 5^4. \quad (7)$$

And this does not match (4). So our noted high-degree vertex having no balancing low-degree vertex suffices to dash any hope of the Pappus graph being in the more distinguished class of the heterocomplementary graphs.

Principle 1 Suppose that

$$E(G; N) > E_{\text{crit}}(N). \quad (8)$$

Then it is usually easier to understand the structure of \overline{G} than that of G . It is usually also more straightforward to name graphs via whichever of themselves or their complement has smaller size (= edge number E). For edge-critical graphs, there is no a priori reason for one to be more suitable than the other. See [60] for some further selection principles in this critical edge case.

⁵The alias *self-complementary* has so far been more widely used.

2.6 A few properties of the Pappus complement




Figure 7:

Remark 10 We unfold row 3 of Fig 6's presentation of $\overline{\text{Pappus}}$ to quickly obtain a 3-crossing presentation in Fig 7.a). And then get it down to 2 crossings in Subfig b).

Subfig d) suffices to demonstrate nonplanarity: it contains, as a subgraph, the utilities graph

$$\text{Utilities} = K_{3,3} :$$

the complete bipartite graph with equal parts of size 3 . Where the second symbol denotes complete-bipartite with 2 pieces of order 3 . By Kuratowski's theorem [40], this is indeed one of the 2 irreducible forbidden subgraphs in a planar graph. The other is the complete graph K_5 .

Exercise 2 Check that Subfig 6.a)'s graph is indeed isomorphic to Subfig 5.e)'s.

3 The Pappus graph is a double irreducible




Figure 8:

Remark 1 $I(\mathfrak{p}_{\text{appus}})$ is cubic and thus has no degree-1 or -2 vertices. Thus it is both foliation irreducible and homeomorph irreducible. Which combination we term double irreducible: class D [54], which we denote with a yellow background, as in Fig 8.

4 Notions of traversability

4.1 The basic notions

Remark 1 The Pappus graph is immediately not *Eulerian*, since it contains odd-degree vertices.

Remark 2 The Pappus graph is straightforwardly *Hamiltonian*.

Remark 3 Consult Chapters 41 and 42 of the freely available [52] if you are not sure what these two notions of traversability mean, or what their basic conceptual content is.

4.2 ZIPHoN treatment

Remark 4 Since the Pappus graph is also planar, one can investigate its Hamiltonian properties using the *ZIPHoN theorem*⁶ By this result, every Hamiltonian cycle splits the graph into 2 outerplanar strips. Each containing an equal amount T of triangulating triangles.

Notational Remark 1 We colour in one outerplanar strip in yellow, leaving the other in white. This includes modelling yellow and white as individually-meaningless and yet mutually-distinguishable labels. By which such strips are invariant under colour exchange. Due to this, one can w.l.o.g. never colour the outer face. We finally mark the bounding Hamiltonian cycles using thick emerald edges.

Remark 5 For the Pappus graph,

$$T = 7.$$

Fig 9 produces 3 such strips, while also indicating some of the forcings by which no more cases are possible.

Remark 6 Our exhaustive case location procedure is to start with a fan of 4 triangles. This fails to give any cases as per row 1. We next consider a continuous fan of 3 triangles, giving the second rows' case. We next consider a contiguous pair, one with a degree-6 vertex and the other not. This

⁶Zero-index planar Hamiltonian necessity theorem [51, 52, 62]. Alias Grinberg's theorem [24, 34, 40].




Figure 9:

is forced either to return to the previous or to give the third row's case. A contiguous pair with both sharing the degree-6 vertex gives our last case. All other cases – 1 or 0 triangles within such a fan – just either become impossible. Or return previous cases, under colour reversal if needs be.

4.3 Manifestly Hamiltonian presentations

Remark 7 Corresponding manifestly-Hamiltonian presentations of the Pappus graph are given in Fig 10.




Figure 10:

Exercise 3- Show that these graphs are isomorphic to each other and to Subfig 1.b)'s graph.

Remark 8 The idea here is to use a manifest regular N -polygon perimeter in place of picking out a non-obviously realized N -cycle in emerald.

Remark 9 We finally make use of both the inner face and the outer face to retain manifest planarity in Fig 11.




Figure 11:

Exercise 4- Use this technique as another way of arriving at a planar presentation for the Fano graph [55], now from its manifestly-Hamiltonian presentation.

5 Notions of colourability




Figure 12:

Remark 1 Since the Pappus graph has a degree-6 vertex, it is ≥ 6 -edge colourable. In fact, it is precisely 6-edge colourable, as per Fig 12.a).

Remark 2 Since the Pappus graph contains triangles, it is ≥ 3 -colourable. And since it is planar, it is ≤ 4 -colourable by the famous 4-colour theorem. In fact, it is 3-colourable, as per Subfig b).

Remark 3 See e.g. [27, 40] or Chapter 47 of the freely-available [52] for an introduction to these notions of colorability. The first two of these are also useful first references for the 4-colour theorem. See [35] for a more advanced account of colourability.

Remark 4 With the Graph Atlas [37] tabulating unique colourability as well, we draw out Subfig c) to demonstrate that the Pappus graph does not have this further property.

Exercise 4⁻ Establish Remark 1 and 2's colourability properties.

6 Pointers to allied material

Pointer 2 A more systematic study would start with Pappus’ law’s simpler configuration. This shall be covered within our article on the simplest Projective graphs [63]. Where no Projective plane conditions, or adjectives, are required. This shall for instance also cover the Pasch configuration [10] and scissor [42] graphs.

Pointer 3 A subsequent systematic study 7 should also cover degenerate cases, such as Pappus’ little theorem. Forming overall the *arena of Pappus configurations*

$$\mathfrak{P}_{\text{Pappus}} .$$

Therein, our optimal subsection’s 1-parameter family forms

$$\mathfrak{Rect}\mathfrak{P}_{\text{Pappus}} :$$

the *subarena of rectangularly-symmetric Pappus configurations*.

Pointer 4 In modelling the arena of Pappus configurations at the level of graphs, a key role is played by the *collineomorphisms* [51]. I.e. the graph isomorphisms that preserve a prescribed set of collinearities: Projectively-significant information. Within this restriction, one cannot access manifestly-planar graph presentations. Leaving our optima as nicest eligible presentations.

Pointer 5 Incidence is the central notion in Projective Geometry. The *incidence graph* for the Pappus configuration shall first be covered in 6; Graph Theorists often call incidence graphs *Levi graphs*. They and Geometers often call the Pappus incidence graph the Pappus graph. Whereas the current Article takes this name to mean the Pappus configuration graph; an older alias for configuration graphs is *Menger graphs* [17].

Pointer 6 Hilbert showed that Pappus’ theorem is equivalent to Algebraic commutativity [11, 42], conferring further Foundational significance.

Pointer 7 The Dandelin–Gallucci property and theorem [49] in $3-d$ are closely related to Pappus’ Theorem. This makes use of the Projective notion of transversality [61] in the context of the generic triple of lines supported in $3-d$.

Pointer 8 The Pappus configuration $\mathfrak{p}_{\text{Pappus}}$ is double-tied in third place [36] for smallest Projective configuration. The smallest is the Fano plane, 7_3 [55]. With a unique proxime 8_3 : Möbius–Kantor’s [17]. And then Pappus arrives, accompanied by 2 other 9_3 ’s. [67] shall more eventually cover the 8_3 and the 2 other 9_3 ’s as graphs; for now see [36, 44, 46] at the level of configurations.

Pointer 9 The *Desargues configuration* is then one of the $10-10_3$ ’s. The corresponding *Desargues’ theorem* [5, 15, 16, 19, 22, 18, 26, 31, 38, 41, 45]. is *the other* fundamental theorem of Projective Geometry. The Desargues configuration, theorem and graph are covered in [58]. Again, this is not to be confused with a more famous incidence graph, which is covered instead in [59].

Non-Desarguian Projective planes [43] are more often mentioned than non-Pappian ones. In fact, for finite planes, Pappus’ theorem implies Desargues’. So conversely in the finite realm, non-Desarguian planes are guaranteed to be non-Pappian. And many Discrete Mathematics ventures confine themselves to finite cases.

Pointer 10 Non-Pappus (Fig 13) is the minimum matroid that is not representable over any field. This can be traced back to all fields enjoying commutativity, which the Pappus matroid encodes but the non-Pappus matroid fails to encode...




Figure 13:

Acknowledgments I thank S. Sánchez and A. Ford for discussions. And the Applied Combinatorics and Topology Discussion Group members.

References

- [1] Menelaus of Alexandria, *Sphaerica* (1st and 2nd Centuries C.E.). See T.A. Heath, *A History of Greek Mathematics. Volume 2, from Aristarchus to Diophantus* (C.U.P., Cambridge, 1921). for a short commentary, or, for a translation and long commentaries, R. Rashed and A. Papadopoulos, *Menelaus' Spherics* (Gruyter, Berlin 2017).
- [2] Pappus of Alexandria, *Synagoge (Collection)* Volume **VII** (4th century C.E.). See in particular A. Jones, *7 of the Collection* (Springer-Verlag 1986) for not only an English Translation but also commentary bridging between the Pappian and modern terminology.
- [3] King Hud of Zaragoza alias al-Mu'taman, *Kitab al-Istikmal (Book of Perfection)* (11th century). 'Ceva's theorem' can already be found here.
- [4] B. Pascal, "Essay pour les Coniques (Essay on Conics)" (1640). For an English translation, see e.g. D.E. Smith, *A Source Book in Mathematics* (McGraw-Hill 1929; Dover, New York 1959).
- [5] G. Desargues' theorem first appeared in print in an Appendix "Manière universelle de M. Desargues pour practiquer la Perspective (Universal Method of Mr. Desargues for using Perspective)" in A. Bosse, *Moyen Universel de Pratiquer la Perspective sur les Tableaux* (1648).
- [6] G. Ceva, *De Lineis Rectis (On Straight Lines)* (1678).
- [7] L.N. Marguerite (Count of Carnot), *Essai sur la Théorie des Transversales (Essay on the Theory of Transversals)* (Courcier, Paris, 1806).
- [8] J-V. Poncelet, *Traité des Propriétés Projectives des Figures (Treatise on the Projective Properties of Figures)* (Bachelier, Paris, 1822).
- [9] K.G.C. von Staudt, *Geometrie der Lage (Geometry of location)* (Verlag, Nürnberg 1847).
- [10] M. Pasch, *Vorlesungen uber neuere Geometrie (Lectures on Modern Geometry)* (Teubner, Leipzig 1882).
- [11] D. Hilbert, *Grundlagen der Geometrie*, (Teubner, Berlin, 1899); for an English translation by E.J. Townsend, see *Foundations of Geometry* (Open Court, 1902; 1950).
- [12] O. Veblen and J.W. Young, *Projective Geometry* (Ginn, Boston 1910).
- [13] C.V. Durell, *Modern Geometry: The Straight Line and Circle* (U. California Libraries, 1920; Macmillan, London 1928).
- [14] R.A. Johnson, *Modern Geometry: an Elementary Treatise on the Geometry of the Triangle and the Circle* (Houghton Mifflin, Boston, MA 1929), republished as *Advanced Euclidean Geometry* (Dover, Mineola, N.Y. 1960).
- [15] J.W. Young, *Projective Geometry* (Open Court, 1931).
- [16] J.A. Todd, *Projective and Analytical Geometry* (Pitman, London 1947).

- [17] H.S.M. Coxeter, "Self-Dual Configurations and Regular Graphs." Bull. Amer. Math. Soc. **56** 413 (1950); reprinted as one of the essays in *The Beauty of Geometry. 12 Essays* (Southern Illinois U.P. 1968; Dover, Mineola N.Y. 1999).
- [18] A. Seidenberg, *Lectures in Projective Geometry* (Van Nostrand, Princeton N.J. 1961; 2012).
- [19] H.S.M. Coxeter, *Introduction to Geometry* (Wiley, New York 1961; 1989).
- [20] Hazel Perfect, *Topics in Geometry* (Pergamon, London 1963).
- [21] D. Pedoe, *An Introduction to Projective Geometry* (Pergamon, Oxford 1963).
- [22] H.S.M. Coxeter, *Projective Geometry* (Blaisdell 1964; 2003).
- [23] H.S.M. Coxeter and S.L. Greitzer, *Geometry Revisited* (M.A.A., Washington, D.C. 1967).
- [24] E.J. Grinberg, "Plane Homogeneous Graphs of Degree Three without Hamiltonian Circuits", Latvian Math. Yearbook **4** 51 (1968); for an English translation, see arXiv:0908.2563 .
- [25] A.S. Posamentier and C.T. Salkind, *Challenging Problems in Geometry* (MacMillan, N.Y. 1970; Dover, N.Y. 1996).
- [26] A.F. Horadam, *A Guide to Undergraduate Projective Geometry* (Pergamon Press , Australia 1970).
- [27] R.J. Wilson, *Introduction to Graph Theory* (Longman, Edinburgh 2010). The first edition dates to 1972.
- [28] A. Seidenberg, "Pappus Implies Desargues" A.M.S. Monthly **83** 192 (1976).
- [29] H.S.M. Coxeter, "The Pappus Configuration and its Groups", Pi Mu Epsilon Journal **6** 331 (1977).
- [30] "The Pappus Configuration and the Self-inscribed Octagon. I.", *Indagationes Mathematicae (Proceedings)* **80** 256 (1977).
- [31] Z.A. Melzak, *Invitation to Geometry* (Wiley, New York 1983).
- [32] B. Grünbaum and G.C. Shephard, *Tilings and Patterns* (Freeman, 1987).
- [33] R. Honsberger, *Episodes in Nineteenth and Twentieth Century Euclidean Geometry* (M.A.A., Washington DC, 1995).
- [34] V.K. Balakrishnan, *Graph Theory* (McGraw–Hill, New York 1997).
- [35] B. Bollobás, *Modern Graph Theory* (Springer, New York 1998).
- [36] B. Polster, *A Geometrical Picture Book* (Springer, New York 1998).
- [37] R.C. Read and R.J. Wilson, *An Atlas of Graphs* (O.U.P., New York 1998).
- [38] V.V. Prasolov and V.M. Tikhomirov, *Geometry* (A.M.S., 2001).
- [39] J.R. Silvester, *Geometry Ancient and Modern* (O.U.P., New York 2001).
- [40] D.B. West, *Introduction to Graph Theory* (Prentice–Hall, Upper Saddle River N.J. 2001).
- [41] M. Reid and B. Szendrői, *Geometry and Topology* (C.U.P. Cambridge 2005).
- [42] J. Stillwell, *The Four Pillars of Geometry* (Springer, New York 2005).
- [43] C. Weibel, "Survey of Non-Desarguesian Planes", Notices of the AMS **54** 1294 (2007).
- [44] B. Grünbaum, *Configurations of Points and Lines* (A.M.S., Providence RI. 2009).
- [45] J. Richter-Gebert, *Perspectives on Projective Geometry: A guide through Real and Complex Geometry* (Springer 2011).
- [46] T. Pisanski and Brigitte Servatius, *Configurations from a Graphical Viewpoint*, (Birkhäuser, Boston 2013).
- [47] I.E. Leonard, J.E. Lewis, A.C.F. Liu and G.W. Tokarsky, *Classical Geometry. Euclidean, Transformational, Inversive and Projective* (Wiley, Hoboken N.J. 2014).
- [48] *Handbook of Graph Drawing and Visualization* ed. R. Tamassia (Taylor & Francis, Boca Raton, Fl. 2014).
- [49] Á.G. Horváth, "Gallucci's Axiom revisited", arXiv.1712.04800

- [50] A. Bogomolny, "Pappus' Theorem" (2018), <https://www.cut-the-knot.org/pythagoras/Pappus.shtml> .
- [51] Combinatorial and Geometrical discussions between E.A. and S. Sánchez (2018).
- [52] E. Anderson, *Applied Combinatorics*, Widely-Applicable Mathematics Series. A. Improving understanding of everything with a pinch of Combinatorics. **0**, (2022). Made freely available in response to the pandemic here: institute-theory-stem.org/combinatorics/
- [53] Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/online-encyclopaedia-of-graphs-and-orders/ .
- [54] E. Anderson and A. Ford (she/her), "Simple Graphs' 8 Double-Irreducibility Classes", Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-graph-double-irreducibility-classes/ (2026).
- [55] E. Anderson, "The Fano Configuration, Plane and Graph", Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-fano/ (2026).
- [56] "The Fano Incidence Graph, alias Heawood Graph and $(6, 3)$ -Cage, Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-fano-incidence-graph/ (2026).
- [57] "The Pappus Incidence Graph", Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-pappus-incidence-graph/ (2026).
- [58] "The Desargues Configuration, Theorem and Graph", Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-desargues-graph/ (2026).
- [59] "The Desargues Incidence Graph", Online Encyclopaedia of Applied Graph and Order Theory, institute-theory-stem.org/oeagot-desargues-incidence-graph/ (2026).
- [60] E. Anderson and A. Ford, *The 6-Graphs*, forthcoming (2026), for [53].
- [61] E. Anderson, *The Structure of Flat Geometry*, forthcoming (2026).
- [62] "Grinberg's Theorem as the Zero-Index Planar Hamiltonian Necessity Theorem. I."; Outerplanar, Triangulation-Count Balance and Zero-Index conceptualizations. forthcoming (2027), for [53].
- [63] "The Smallest Projective Diagrams and their Graphs", forthcoming (2027), for [53].
- [64] *The Structure of Differential Geometry, Widely-Applicable Mathematics. A. Improving understanding of everything by dual-wielding Combinatorics and Algebra. 6*, forthcoming 2028.
- [65] *Lie Algebras and their Representations*, Widely-Applicable Mathematics Series A. Improving understanding of everything by dual-wielding Algebra and Combinatorics **7**, forthcoming 2029.
- [66] "Variations on Pappus' Configuration, Theorem, and Graphs", forthcoming (2030) for [53].
- [67] "The remaining Smallest Projective Configurations and their Graphs", forthcoming (2030) for [53]. An account of Möbius–Kantor is however expected to appear in 2027...